

Section 3.8

Derivative of the inverse function and logarithms

3 Lecture

Dr. Abdulla Eid

College of Science

MATHS 101: Calculus I

- 1 Inverse Functions (1 lecture).
- 2 Logarithms.
- 3 Derivative of the inverse function (1 lecture).
- 4 Logarithmic differentiation (1 lecture).

Dr. Abdulla Eid

1 - Inverse functions (pre-calculus)

Definition

Let f be a function. The **inverse** function, denoted by f^{-1} of f is a *new* function such that

$$\underbrace{f}_{\text{outer}}(\underbrace{f^{-1}}_{\text{inner}}(x)) = x \quad \text{and} \quad \underbrace{f^{-1}}_{\text{outer}}(\underbrace{f}_{\text{inner}}(x)) = x$$

(The function and its inverse cancel each other).

Example

(a) Let $f(x) = x + 5$, then $f^{-1}(x) = x - 5$ (we will see how to find the inverse shortly). Then

- $f(f^{-1}(x)) = f(x - 5) = x - 5 + 5 = x.$
- $f^{-1}(f(x)) = f^{-1}(x + 5) = x + 5 - 5 = x.$

(b) Let $f(x) = x^2 (x \geq 0)$, then $f^{-1}(x) = \sqrt{x}$ because:

- $f(f^{-1}(x)) = f(\sqrt{x}) = (\sqrt{x})^2 = x.$
- $f^{-1}(f(x)) = f^{-1}(x^2) = \sqrt{x^2} = |x| = x.$

Question: has every function an inverse? How to tell when a function has an inverse?

Answer: No, we use the **horizontal line test** if we have the graph of the function.

To find the inverse function

To find the inverse function

Algebraically

Step 1: Write $y = f(x)$.

Step 2: Switch x
and y to get $x = f(y)$.

Step 3: Solve for y ,
i.e., isolate y alone to get
 $y = f^{-1}(x)$.

Geometrically

Step 1: Reflect the graph of $y = f(x)$
on the x -axis.

Step 2: rotate the resulting graph
by 90° counterclockwise to get the graph
of $f^{-1}(x)$.

Example

Find the inverse of $g(x) = 5x - 3$.

Solution:

Step 1: Write $y = g(x) \rightarrow y = 5x - 3$.

Step 2: Exchange x and y in step 1 $\rightarrow x = 5y - 3$.

Step 3: Solve the equation in step 1 for y

$$x = 5y - 3$$

$$x + 3 = 5y$$

$$\frac{x + 3}{5} = y$$

Hence we have

$$g^{-1}(x) = \frac{x + 3}{5}$$

Exercise

Find the inverse function of

① $f(x) = 3x + 2.$


② $f(x) = x^2 - 1 (x > 0).$

③ $f(x) = \frac{1}{x}.$

④ $f(x) = \sqrt{x}.$


Example

Find the graph of the inverse function of the following functions:




Example

Find the graph of the inverse function of the following functions:



Example

Find the graph of the inverse function of the following functions:



Inverse Trigonometric Functions

Example

Let $y = f(x) = \sin x$. Then the graph of the $f(x)$ is given by

Therefore, f has an inverse if $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ and we write it as

$$f^{-1}(x) = \sin^{-1} x = \arcsin x.$$

- 1 Domain of \sin^{-1} is $[-1, 1]$.
- 2 Range of \sin^{-1} is $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

Inverse Trigonometric Functions

Example

Let $y = f(x) = \cos x$. Then the graph of the $f(x)$ is given by

Therefore, f has an inverse if $x \in [0, \pi]$ and we write it as

$$f^{-1}(x) = \cos^{-1} x = \arccos x.$$

- 1 Domain of \cos^{-1} is $[-1, 1]$.
- 2 Range of \cos^{-1} is $[0, \pi]$.

Inverse Trigonometric Functions

Example

Let $y = f(x) = \tan x$. Then the graph of the $f(x)$ is given by

Therefore, f has an inverse if $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ and we write it as

$$f^{-1}(x) = \tan^{-1} x = \arctan x.$$

- 1 Domain of \tan^{-1} is $[-\infty, \infty$.
- 2 Range of \tan^{-1} is $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.

Exercise

Find the domain, range, and the graph of inverse of the following functions:

① $f(x) = \cot x.$

② $f(x) = \sec x.$

③ $f(x) = \csc x.$

Dr. Abdulla